

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal621con

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 DEC 01 ChemPort single article sales feature unavailable
NEWS 3 APR 03 CAS coverage of exemplified prophetic substances enhanced
NEWS 4 APR 07 STN is raising the limits on saved answers
NEWS 5 APR 24 CA/CAPLUS now has more comprehensive patent assignee information
NEWS 6 APR 26 USPATFULL and USPAT2 enhanced with patent assignment/reassignment information
NEWS 7 APR 28 CAS patent authority coverage expanded
NEWS 8 APR 28 ENCOMPLIT/ENCOMPLIT2 search fields enhanced
NEWS 9 APR 28 Limits doubled for structure searching in CAS REGISTRY
NEWS 10 MAY 08 STN Express, Version 8.4, now available
NEWS 11 MAY 11 STN on the Web enhanced
NEWS 12 MAY 11 BEILSTEIN substance information now available on STN Easy
NEWS 13 MAY 14 DGENE, PCTGEN and USGENE enhanced with increased limits for exact sequence match searches and introduction of free HIT display format
NEWS 14 MAY 15 INPADOCDB and INPAFAMDB enhanced with Chinese legal status data
NEWS 15 MAY 28 CAS databases on STN enhanced with NANO super role in records back to 1992
NEWS 16 JUN 01 CAS REGISTRY Source of Registration (SR) searching enhanced on STN
NEWS 17 JUN 26 NUTRACEUT and PHARMAML no longer updated
NEWS 18 JUN 29 IMSCOPROFILE now reloaded monthly
NEWS 19 JUN 29 EPFULL adds Simultaneous Left and Right Truncation (SLART) to AB, MCLM, and TI fields
NEWS 20 JUL 09 PATDPAFULL adds Simultaneous Left and Right Truncation (SLART) to AB, CLM, MCLM, and TI fields
NEWS 21 JUL 14 USGENE enhances coverage of patent sequence location (PSL) data
NEWS 22 JUL 27 CA/CAPLUS enhanced with new citing references
NEWS 23 JUL 16 GBFULL adds patent backfile data to 1855
NEWS 24 JUL 21 USGENE adds bibliographic and sequence information
NEWS 25 JUL 28 EPFULL adds first-page images and applicant-cited references
NEWS 26 JUL 28 INPADOCDB and INPAFAMDB add Russian legal status data

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:04:05 ON 01 AUG 2009

=> FILE CASREACT

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.44

0.44

FILE 'CASREACT' ENTERED AT 15:04:57 ON 01 AUG 2009

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 26 Jul 2009 VOL 151 ISS 5

New CAS Information Use Policies, enter HELP USAGETERMS for details.


* CASREACT now has more than 16.5 million reactions *
* *

CASREACT contains reactions from CAS and from: ZIC/VINITI database (1974-1999) provided by InfoChem; INPI data prior to 1986; Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich; organic reactions, portions copyright 1996-2006 John Wiley & Sons, Ltd., John Wiley and Sons, Inc., Organic Reactions Inc., and Organic Syntheses Inc. Reproduced under license. All Rights Reserved.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=>

Uploading C:\Program Files\Stnexp\Queries\GGSS64.str



```

chain nodes :
1  2  3  4  5  6  7  8  9  10  11  12  13  14  15  16  17  18  19  20  21  22
chain bonds :
1-2  1-8  1-9  2-3  2-10  3-4  3-11  5-6  6-7  12-13  12-17  13-14  13-21  13-22
14-15  14-19  14-20  15-16  15-18
exact/norm bonds :
3-11  12-13  15-18
exact bonds :
1-2  1-8  1-9  2-3  2-10  3-4  5-6  6-7  12-17  13-14  13-21  13-22  14-15  14-19
14-20  15-16

```

```

Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
18:CLASS 19:CLASS 20:CLASS 21:CLASS 22:CLASS
fragments assigned product role:
containing 12
fragments assigned reactant/reagent role:
containing 1
containing 5

```

L1 STRUCTURE UPLOADED

=> S L1 FULL

FULL SEARCH INITIATED 15:05:28 FILE 'CASREACT'

SCREENING COMPLETE - 47118 REACTIONS TO VERIFY FROM 4466 DOCUMENTS

100.0% DONE 47118 VERIFIED

13 HIT RXNS

10 DOCS

SEARCH TIME: 00.00.08

L2 10 SEA SSS FUL L1 (13 REACTIONS)

=> S L2 ABD BASE

MISSING OPERATOR L2 ABD

The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> S L2 AND BASE AND ACID

41867 BASE

255988 ACID

L3 1 L2 AND BASE AND ACID

=> D L3 IBIB ABS CRD

L3 ANSWER 1 OF 1 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER:

140:356948 CASREACT

TITLE:

Catalytic addition reaction for the production of 3-(methylthio)propanal from mercaptomethane and acrolein

INVENTOR(S):

Rey, Patrick

PATENT ASSIGNEE(S):

Adisseo France S.A.S., Fr.

SOURCE:

Eur. Pat. Appl., 10 pp.

CODEN: EPXXDW

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1413573	A1	20040428	EP 2002-356211	20021024
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
CA 2495746	A1	20040506	CA 2003-2495746	20031014
WO 2004037774	A1	20040506	WO 2003-IB4557	20031014
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

AU 2003267771 EP 1556343 EP 1556343 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK BR 2003015385 CN 1705641 CN 1277816 JP 2006515834 AT 371642 ES 2291662 RU 2336266 ZA 2005001389 MX 2005004158 US 20050240048 US 7256315 NO 2005002471 A1 20040513 A1 20050727 B1 20070829 A 20050823 A 20051207 C 20061004 T 20060608 T 20070915 T3 20080301 C2 20081020 A 20060726 A 20050803 A1 20051027 B2 20070814 A 20050725	AU 2003-267771 EP 2003-748466 BR 2003-15385 CN 2003-80101589 JP 2004-546263 AT 2003-748466 ES 2003-748466 RU 2005-105040 ZA 2005-1389 MX 2005-4158 US 2005-524548 NO 2005-2471 EP 2002-356211 WO 2003-1B4557	20031014 20031014 20031014 20031014 20031014 20031014 20050216 20050419 20050516 20050523 20021024 20031014
--	---	--

PRIORITY APPLN. INFO.:

AB A process for the production of 3-(methylthio)propanal comprises reacting mercaptomethane and acrolein in the presence of a catalyst comprising an organic base such as an N-alkylmorpholine (e.g., 4-methylmorpholine).

RX(1) OF 3



NOTE: optimization study, optimized on catalyst

CON: STAGE(1) room temperature -> 40 deg C; 40 deg C

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> D L2 IBIB ABS CRD 1-10

L2 ANSWER 1 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER:

149:448709 CASREACT

TITLE:

Synthesis of methionine- and norleucine-derived phosphinopeptides

AUTHOR(S):

Liboska, Radek; Picha, Jan; Hanclova, Ivona;
 Budesinsky, Milos; Sanda, Miloslav; Jiracek, Jiri
 Institute of Organic Chemistry and Biochemistry,
 Academy of Sciences of the Czech Republic, Prague 6,
 166 10, Czech Rep.

SOURCE:

Tetrahedron Letters (2008), 49(39), 5629-5631

CODEN: TELEAY; ISSN: 0040-4039

PUBLISHER:

Elsevier Ltd.

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB We present a straightforward synthesis of N-Fmoc-protected synthons derived from a phosphinic analog of methionine. These precursors were used successfully for the solid-phase synthesis of methionine-mimic phosphinopeptides using BOP-catalyzed coupling without protection of the

phosphoryl moiety. We also prepared a new type of pseudopeptide derived from a phosphinic analog of norleucine with a -PO(OH)CH₂CO₂R moiety.

RX(1) OF 110



NOTE: Michael addition
CON: 0 deg C

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 2 OF 10 CASREACT COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 144:214741 CASREACT
TITLE: Method and catalysts for preparing
3-(methylthio)propanal from acrolein and methyl
mercaptan and for the manufacture of
2-hydroxy-4-(methylthio)butanenitrile from it and
hydrogen cyanide
INVENTOR(S): Dubner, Frank; Weckbecker, Christoph
PATENT ASSIGNEE(S): Germany
SOURCE: U.S. Pat. Appl. Publ., 8 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20060030739	A1	20060209	US 2005-198609	20050805
US 7119233	B2	20061010		
DE 102004038053	A1	20060427	DE 2004-10200403805320040805	
CA 2573047	A1	20060216	CA 2005-2573047	20050714
WO 2006015684	A2	20060216	WO 2005-EP7666	20050714
WO 2006015684	A3	20060803		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
EP 1778631	A2	20070502	EP 2005-775924	20050714
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR			
CN 101031542	A	20070905	CN 2005-80026591	20050714
JP 2008508330	T	20080321	JP 2007-524209	20050714
BR 2005013072	A	20080422	BR 2005-13072	20050714

MX 2007000345 A 20070307 MX 2007-345 20070109
IN 2007KN00106 A 20070629 IN 2007-KN106 20070109
PRIORITY APPLN. INFO.: DE 2004-10200403805320040805
WO 2005-EP7666 20050714

OTHER SOURCE(S): MARPAT 144:214741

AB A method is described for preparing 3-(methylthio)propanal (I) by the the addition reaction of Me mercaptan to acrolein in the presence of macro-reticular resin catalysts containing pendant tertiary-amine groups [e.g., [(dimethylamino)methyl]styrene copolymer] to give I which is then reacted with HCN in the presence of the same catalyst to give 2-hydroxy-4-(methylthio)butanenitrile. Process flow diagrams are presented.


RX(1) OF 7



NOTE: solid-supported catalyst on Merrifield resin,
3-(methylthio)propanal used as reaction medium, batchwise
synthesis

CON: STAGE(1) 10 minutes, 0 deg C
STAGE(2) 2 hours, 0 deg C


RX(3) OF 7



NOTE: solid-supported catalyst on Merrifield resin,
3-(methylthio)propanal used as reaction medium, continuous
synthesis

CON: STAGE(1) 30 minutes, 50 deg C; 30 minutes, 40 deg C


RX(4) OF 7



NOTE: solid-supported catalyst on Merrifield resin,
3-(methylthio)propanal used as reaction medium, batchwise
synthesis

CON: STAGE(1) 10 minutes, 0 deg C
STAGE(2) 2 hours, 0 deg C

RX(6) OF 7



NOTE: solid-supported catalyst on Merrifield resin,
3-(methylthio)propanal used as reaction medium, continuous
synthesis

CON: STAGE(1) 30 minutes, 50 deg C; 30 minutes, 40 deg C

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 3 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 140:356948 CASREACT

TITLE: Catalytic addition reaction for the production of 3-(methylthio)propanal from mercaptomethane and acrolein

INVENTOR(S): Rey, Patrick

PATENT ASSIGNEE(S): Adisseo France S.A.S., Fr.

SOURCE: Eur. Pat. Appl., 10 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1413573	A1	20040428	EP 2002-356211	20021024
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
CA 2495746	A1	20040506	CA 2003-2495746	20031014
WO 2004037774	A1	20040506	WO 2003-1B4557	20031014
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003267771	A1	20040513	AU 2003-267771	20031014
EP 1556343	A1	20050727	EP 2003-748466	20031014
EP 1556343	B1	20070829		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003015385	A	20050823	BR 2003-15385	20031014
CN 1705641	A	20051207	CN 2003-80101589	20031014
CN 1277816	C	20061004		
JP 2006515834	T	20060608	JP 2004-546263	20031014
AT 371642	T	20070915	AT 2003-748466	20031014
ES 2291662	T3	20080301	ES 2003-748466	20031014
RU 2336266	C2	20081020	RU 2005-105040	20031014
ZA 2005001389	A	20060726	ZA 2005-1389	20050216
MX 2005004158	A	20050803	MX 2005-4158	20050419
US 20050240048	A1	20051027	US 2005-524548	20050516
US 7256315	B2	20070814		
NO 2005002471	A	20050725	NO 2005-2471	20050523
PRIORITY APPLN. INFO.:			EP 2002-356211	20021024
			WO 2003-1B4557	20031014
AB A process for the production of 3-(methylthio)propanal comprises reacting mercaptomethane and acrolein in the presence of a catalyst comprising an				

organic base such as an N-alkylmorpholine (e.g., 4-methylmorpholine).

RX(1) OF 3



NOTE: optimization study, optimized on catalyst

CON: STAGE(1) room temperature -> 40 deg C; 40 deg C

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 4 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 140:287102 CASREACT
TITLE: Method for producing 3-methylthiopropional from acrolein and methyl mercaptan

INVENTOR(S): Shiozaki, Tetsuya; Haga, Toru
PATENT ASSIGNEE(S): Sumitomo Chemical Company, Limited, Japan
SOURCE: U.S. Pat. Appl. Publ., 4 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent
LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040063650	A1	20040401	US 2003-665006	20030922
JP 2004115461	A	20040415	JP 2002-282874	20020927
JP 4186572	B2	20081126		
EP 1408029	A1	20040414	EP 2003-21191	20030924
EP 1408029	B1	20061122		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
ES 2275989	T3	20070616	ES 2003-21191	20030924
CN 1496979	A	20040519	CN 2003-125534	20030925
CN 100349863	C	20071121		
IN 2003CH00786	A	20051118	IN 2003-CH786	20030925
			JP 2002-282874	20020927

PRIORITY APPLN. INFO.:

AB 3-Methylthiopropional is produced in high yield and selectivity by supplying acrolein and Me mercaptan together or sequentially with an acidic compound (e.g., acetic acid) and a basic compound (e.g., pyridine) into a reaction system to react the acrolein with the Me mercaptan, where the basic compound is used in an amount of about 0.3 mol or less per mol of the acidic compound

RX(1) OF 1



NOTE: other products detected

CON: 45 - 50 minutes, 70 deg C

L2 ANSWER 5 OF 10 CASREACT COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 128:114715 CASREACT
 TITLE: Processes for the preparation of
 3-(methylthio)propanal and
 2-hydroxy-4-(methylthio)butanenitrile
 INVENTOR(S): Blackburn, Thomas F.; Pellegrin, Paul F.
 PATENT ASSIGNEE(S): Novus International, Inc., USA
 SOURCE: U.S., 9 pp., Cont.-in-part of U.S. 5,663,409.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5705675	A	19980106	US 1995-581249	19951229
US 5663409	A	19970902	US 1995-476356	19950607
ZA 9604335	A	19960820	ZA 1996-4335	19960528
WO 9640631	A1	19961219	WO 1996-US9060	19960604
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML				
AU 9659873	A	19961230	AU 1996-59873	19960604
AU 714151	B2	19991223		
EP 830341	A1	19980325	EP 1996-917222	19960604
EP 830341	B1	20010905		
R: BE, DE, DK, ES, FR, GB, IT, LU, NL, MC, PT, IE				
CN 1189818	A	19980805	CN 1996-195190	19960604
CN 1092184	C	20021009		
JP 11511119	T	19990928	JP 1997-501471	19960604
RU 2173681	C2	20010920	RU 1998-100220	19960604
ES 2160819	T3	20011116	ES 1996-917222	19960604
PT 830341	T	20011228	PT 1996-917222	19960604
CN 1510030	A	20040707	CN 2002-2002126457	19960604
PRIORITY APPLN. INFO.:			US 1995-476356	19950607
			US 1995-581249	19951229
			WO 1996-US9060	19960604

OTHER SOURCE(S): MARPAT 128:114715

AB A catalytic processes for the preparation of 3-(methylthio)propanal and 2-hydroxy-4-(methylthio)butanenitrile using novel addition catalysts is described. The novel addition catalysts include: triisopropanolamine, nicotinamide, imidazole, benzimidazole, 2-fluoropyridine, poly-4-vinylpyridine, 4-dimethylaminopyridine, picoline, pyrazine, trialkylamines, and tertiary amines. E.g., reaction of MeSH and acrolein in presence of poly-4-vinylpyridine gave 89.0% 3-(methylthio)propanal. The aldehyde product, containing the poly-4-vinylpyridine catalyst, was converted to the nitrile in the same reactor by treatment with HCN. The yield of nitrile was 72.9%.

RX(1) OF 3



NOTE: novel process focuses on the catalyst/acid combination; process minimizes the extent of polymer formation

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 6 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 126:157183 CASREACT

TITLE: Process for the continuous preparation of 3-(methylthio)propanal from acrolein and methyl mercaptan

INVENTOR(S): Hsu, Yung C.

PATENT ASSIGNEE(S): Novus International, Inc., USA

SOURCE: PCT Int. Appl., 85 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9700858	A1	19970109	WO 1996-US10920	19960621
W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML			
US 5905171	A	19990518	US 1996-667099	19960620
AU 9663959	A	19970122	AU 1996-63959	19960621
AU 726921	B2	20001123		
EP 842149	A1	19980520	EP 1996-923452	19960621
EP 842149	B1	20030205		
R:	BE, DE, DK, ES, FR, GB, IT, LU, NL, MC, PT, IE			
CN 1188470	A	19980722	CN 1996-194943	19960621
CN 1120834	C	20030910		
JP 11508266	T	19990721	JP 1997-504005	19960621
RU 2172734	C2	20010827	RU 1998-100590	19960621
ES 2192607	T3	20031016	ES 1996-923452	19960621
PRIORITY APPLN. INFO.:			US 1995-421P	19950622
			US 1996-667099	19960620
			WO 1996-US10920	19960621

AB In the title process, a liquid reaction, medium containing 3-(methylthio)propanal and a catalyst for the reaction between Me mercaptan and acrolein, is contacted with a gaseous acrolein feed stream in a gas-liquid contact zone. The gaseous acrolein feed stream comprises acrolein vapor and noncondensable gas and the acrolein is transferred from the acrolein feed stream to the reaction medium. Me mercaptan, introduced into the reaction medium, reacts with the acrolein in that medium,

producing a liquid reaction product containing 3-(methylthio)propanal. The noncondensable gas is then separated from the liquid reaction product the reaction product is divided into a produce fraction and a circulating fraction, and the circulating fraction is recycled to the gas/liquid contact zone. Process flow diagrams are presented.

RX(1) OF 1



NOTE: continous process


REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 7 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 124:184625 CASREACT
 TITLE: Process for the treatment and conditioning of solid or liquid effluents charged with heavy metals
 INVENTOR(S): Leybros, Jean
 PATENT ASSIGNEE(S): Commissariat a l'Energie Atomique, Fr.
 SOURCE: Eur. Pat. Appl., 9 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 687483	A1	19951220	EP 1995-401367	19950613
EP 687483	B1	19980826		
R: BE, CH, DE, ES, GB, IT, LI, NL				
FR 2721237	A1	19951222	FR 1994-7297	19940615
FR 2721237	B1	19960802		
ES 2123221	T3	19990101	ES 1995-401367	19950613
PRIORITY APPLN. INFO.:				
AB	The effluent is treated with a reducing agent (e.g., SO ₂) and then contacted with an organic extractant (e.g., bis(2-ethylhexyl)phosphoric acid) and a hydrocarbon (e.g., hydrogenated tetrapropylene) for selective removal of the metal ions, followed by removing the heavy metals from the organic extract by a 2nd aqueous extraction, and precipitating and filtering the metals from the aqueous solution			

RX(1) OF 1



NOTE: Classification: S-Alkylation; "1,4-Addition"; # Conditions: (AcO)₂; <50 deg 2atm; # Comments: 4.7.49


L2 ANSWER 8 OF 10 CASREACT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 120:133858 CASREACT
 TITLE: Process for producing 2-hydroxy-4-methylthiobutanoic acid
 INVENTOR(S): Matsuoaka, Kazuyuki
 PATENT ASSIGNEE(S): Daicel Chemical Industries, Ltd., Japan
 SOURCE: PCT Int. Appl., 21 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9323372	A1	19931125	WO 1993-JP659	19930520
W: US				
RW: BE, DE, FR, GB				
JP 06049020	A	19940222	JP 1993-143026	19930520
JP 3219544	B2	20011015		
EP 601195	A1	19940615	EP 1993-910360	19930520
EP 601195	B1	19960828		
R: BE, DE, FR, GB				
CN 1084511	A	19940330	CN 1993-107598	19930521
CN 1036391	C	19971112		
US 5386056	A	19950131	US 1994-178315	19940112
PRIORITY APPLN. INFO.:				
			JP 1992-155802	19920521
			WO 1993-JP659	19930520

AB A process for producing 2-hydroxy-4-methylthiobutanoic acid (I) together with methanol comprises hydrating 2-hydroxy-4-methylthiobutyronitrile (II) into 2-hydroxy-4-methylthiobutanamide (III), reacting the amide with Me formate to yield Me 2-hydroxy-4-methylthiobutanoate (IV) and formamide, and hydrolyzing the Me ester. The discharge of a large amount of ammonium sulfate can be prevented, because no sulfuric acid is used as the reactant. The byproduct formamide and methanol are utilizable as the starting material of the reaction after converting them into HCN and Me formate, resp. Thus, addition of MeSH to acrolein in the presence of Cu(OAc)₂ and hydroquinone and addition of the resulting 3-methylthiopropionaldehyde with HCN in the presence of NaOH in MeOH gave II. Hydration of II in the presence of MnO₂ in aqueous acetone at 60° for 6 h to give III which was reacted with HCO₂Me in MeOH containing MeONa to give IV and the byproduct formamide. Hydrolysis of IV in the presence of Amberlyst 15 in H₂O at 95° gave I, while the byproduct MeOH was recovered. Formamide was fed into a stainless steel reactor packed with alumina at 500° to give HCN. MeOH was contacted with a catalyst prepared from Cu(NO₃)₂ and ammonium chromate in a stainless steel reactor to give Me formate.

RX(2) OF 15



NOTE: 20.degree.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 9 OF 10 CASREACT COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 51:47157 CASREACT
TITLE: 3-(Methylthio)propanal
INVENTOR(S): Hunt, Madison; Merner, Richard R.
PATENT ASSIGNEE(S): E. I. du Pont de Nemours & Co.
DOCUMENT TYPE: Patent
LANGUAGE: Unavailable
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	US 2776996		19570108	US 1955-461955	19551222
AB	A mixture of MeSH (I) 440 and pyridine 16 is fed into acrolein 500 and HOAc 5 parts in an autoclave below 75°. The final portion of 3-(methylthio)-propanal (II) and I is added rapidly at 40° to give 91-7% II.				

RX(1) OF 1


$\text{H}_3\text{C}-\text{SH} \xrightarrow[\text{AcOH}]{\text{H}_2\text{C:CHCHO, Pyridine,}} \text{MeS}-\text{CH}_2-\text{CH}_2-\text{CHO}$

NOTE: Classification: S-Alkylation; "1,4-Addition"; # Conditions: MeSH pyridine AcOH; 70-75 deg; # Comments: high yield

L2 ANSWER 10 OF 10 CASREACT COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 42:25284 CASREACT
TITLE: Synthesis of DL-methionine
AUTHOR(S): Pierson, Earl; Giella, Mario; Tishler, Max
CORPORATE SOURCE: Merck & Co., Inc., Rahway, NJ
SOURCE: Journal of the American Chemical Society (1948), 70, 1450-1
CODEN: JACSAT; ISSN: 0002-7863
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable
AB Addition of 48 g. MeSH to 56 g. CH_2CHCHO and 0.5 g. $\text{Cu}(\text{OAc})_2$ at 35-40° gives 84% $\text{MeSCH}_2\text{CH}_2\text{CHO}$ (I), b11 52-4°, n_D20 1.4850, d₂₀ 1.036 (2,4-dinitrophenylhydrazones, m. 116-19°). I (10.4 g.), shaken with 10.4 g. NaHSO_3 in 35 mL. H_2O , the product treated (in 3 portions) with 4.9 g. NaCN in 15 mL. H_2O (temperature below 35°), the oil immediately extracted with C_6H_6 , and the C_6H_6 extracted with NaHSO_3 , gives 90% α -hydroxy- β -(methylmercapto)butyronitrile (II), an oil that distilled at 100°/3 μ . I (26 g.), 113 g. $(\text{NH}_4)_2\text{CO}_3$, 24.5 g. NaCN , 335 mL. EtOH , and 335 mL. H_2O , heated 4 h. at 50-5°, and the filtrate concentrated to 300 mL. and heated 5 min. at 90° with 50 mL. concentrated HCl , give 79% 5-(2-methylmercaptoethyl)hydantoin (III), m. 103-5°; it results in 50% yield (based on I) from II and $(\text{NH}_4)_2\text{CO}_3$ in 50% MeOH (2.5 h. at 50-5°). III (17.4 g.) and 8.8 g. NaOH in 75 mL. H_2O , refluxed 6 h., an addnl. 4.4 g. NaOH added, and the refluxing continued for 18 h., give 10.6 g. DL-methionine (IV), m. 269°

(decomposition); if I and III are not isolated, the yield (based on CH₂:CHCHO) is 50%. II (123 g.), treated 30 min. at 60° with NH₃, gives 40% of crude methionine nitrile, which could not be purified; hydrolysis by heating 5.5 h. on the steam bath with 20 mL. concentrated HCl yields 75% IV. Hydrolysis of III to IV was also effected by concentrated HCl at 135° and by (NH₄)₂S at 135°.

RX(1) OF 1



NOTE: Classification: "1,4-Addition"; S-Alkylation; # Conditions:
Cu(OAc)₂ MeSH gas; 30mn 40 deg; 1h

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	197.58	198.02
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-8.58	-8.58

STN INTERNATIONAL LOGOFF AT 15:07:59 ON 01 AUG 2009